arXiv:math/0008194v1 [math.DS] 25 Aug 2000

ERGODICITY OF MAPPING CLASS GROUP ACTIONS ON REPRESENTATION VARIETIES, II. SURFACES WITH BOUNDARY

DOUG PICKRELL AND EUGENE Z. XIA

Abstract. The mapping class group of a compact oriented surface of genus greater than one with boundary acts ergodically on connected components of the representation variety corresponding to a connected compact Lie group, for every choice of conjugacy class boundary condition.

1. Introduction and Results

Let Σ be a compact oriented surface with $n+1$ boundary components (circles) $(c_0, ..., c_n)$ and K a compact connected Lie group of dimension d and rank r. Denote by Γ the mapping class group of Σ that fixes the $c_j's$. Fix a set of conjugacy classes $\{C_j\}_{j=0}^{n}$, one for each boundary component and let

$$C = \prod_{j=0}^{n} C_j.$$

Then the representation variety is

$$\mathrm{Hom}_C(\pi_1(\Sigma), K) = \{\rho : \rho(c_j) \in C_j\}.$$

The group K acts on $\mathrm{Hom}_C(\pi_1(\Sigma), K)$ by conjugation and the moduli space is the quotient

$$\mathrm{Hom}_C(\pi_1(\Sigma), K)/conj(K).$$

Theorem 1.1. *Fix $p > 1$. With respect to the Lebesgue class, Γ acts ergodically on the connected components of $\mathrm{Hom}_C(\pi_1(\Sigma), K)/conj(K)$.*

When $n = 0$, this theorem was proved in [3].

2. The representation variety and the mapping class group

With slight modifications, the basic notations are as in [3]. Let $K' = [K, K]$ be the semi-simple part of K and dg', dg their respective Haar measures. Denote by $\mathfrak{k}, \mathfrak{k}'$ the Lie algebras of K, K', respectively.

Date: February 1, 2008.

As Σ has boundary, by adding a possible extra boundary component and a fixed central element for that boundary component, one may assume K' is simply connected without losing generality.



FIGURE 1: The surface Σ.

Fix the base point on c_0. Then $\pi_1(\Sigma)$ has the following presentation:

$$\langle x_i, y_i, c_j \;\mid\; \prod_{i=1}^{p}[x_i, y_i]\prod_{j=0}^{n} c_j\rangle.$$

In terms of Figure 1, x_1, y_1 belong to the one-holed torus bounded by s while for $i > 1$, x_i, y_i belong to the one-holed genus $(p-1)$ surface bounded by s'. All the boundary circles in Figure 1 are oriented counter-clockwise with the exception of c_0. The following notions are fixed through out this paper. By a slight abuse of notation, the $c_j's$ will also denote elements in K. Let $g = (g_1, ..., g_p), h = (h_1, ..., h_p) \in K^p$ and $c = (c_0, ..., c_n) \in K^{n+1}$, although in many cases c will denote elements in C. Define maps

$$R_{l,k}(g,h) = \prod_{i=l}^{k} [g_i, h_i],$$

$$S_{l,k}(c) = \prod_{i=l}^{k} c_j.$$

By Remark (2.1.5) in [3], the image of the map $R_{l,k} : K^{2(k-l)} \longrightarrow K'$ has full measure in K'. Since the image of $R_{l,k}$ is compact, $R_{l,k}$ is actually onto K'.

The mapping class group Γ is the group of connected components of the diffeomorphism group of Σ fixing the boundary $\partial\Sigma$. The group Γ is generated by the Dehn twists. In particular, Γ contains the twists along x_i, y_i and σ_j portrayed in Figure 1. Let Γ_0 be the subgroup generated by the Dehn twists along the x_i's and y_i's and Γ_j be the subgroup generated by $\Gamma_0 \cup \{\sigma_j\}$.

3. ERGODICITY

Let $p > 1$. Fix conjugacy classes C_j for each, $0 \leq j \leq n$. Then the representation variety $\mathrm{Hom}_C(\Sigma, K)$ is

$$\{(g,h,c) \in K^{2p} \times C : R_{1,p}(g,h) S_{0,n}(c) = I\}.$$

Assume $\mathrm{Hom}_C(\Sigma, K)$ is non-empty which means that $S_{0,n}(c) \in K'$. Each conjugacy class C_j possess a K-invariant canonical measure. From these measures on the conjugacy classes and the measure dg on K, the variety $\mathrm{Hom}_C(\Sigma, K)$ inherits a Lebesgue measure whose push-forward measure on $\mathrm{Hom}_C(\pi_1(\Sigma), K)/conj(K)$ is in the same measure class as μ.

Fix j and for each $l \neq j$, fix an element $c_l \in C_l$. Let $k = S_{l+1,n}(c)$, $s = R_{1,1}(g,h)$ and $s' = R_{2,p}(g,h)$. Then the action of σ_j fixes c_l for all $j \neq l$ (See Figure 1) and $\sigma_j(c_j) = (skc_jk^{-1})c_j(skc_jk^{-1})^{-1}$. Let

$$M = \{(g,h,c_j) : R_{1,p}(g,h) S_{0,n}(c) = I\} \subset \mathrm{Hom}_C(\pi_1(\Sigma), K).$$

For each fixed $c_j \in C_j$, let

$$N(c_j) = \{(g,h) : R_{1,p}(g,h) S_{0,n}(c) = I\} \subset M.$$

Then both M and $N(c_j)$ inherit measures from $\mathrm{Hom}_C(\Sigma, K)$. Indeed, M and $N(c_j)$ are themselves both representation varieties on a surface with two boundary components. In the case of M, one boundary is associated with a fixed group element and the other a conjugacy class while both boundary components of $N(c_j)$ are associated with group elements. In particular, as subspaces of $\mathrm{Hom}_C(\Sigma, K)$, M is Γ_j-invariant while $N(c_j)$ is Γ_0-invariant.

Let $f \in L^2(M)$ be Γ_j-invariant. The goal is to show that f is constant a.e. The following theorem is from [3]:

Theorem 3.1. *The group Γ_0 acts ergodically on $N(c_j)$.*

This shows that f is constant on $N(c_j)$ for a.e. $c_j \in C_j$. Hence f is a characteristic function on C_j. It remains to show that f is independent of c_j. Consider the function

$$\phi : M \longrightarrow C_j \times C_j :$$

$$\phi(g, h, c_j) = (c_j, (skc_jk^{-1})c_j(skc_jk^{-1})^{-1}).$$

Since f is Γ_j-invariant, to show that f is independent of c_j, it is sufficient to show that the relation defined by $Im(\phi)$ is transitive; since the equality holds in an a.e. sense, it is sufficient to show that

Lemma 3.2. *Let $pr_1 : C_j \times C_j \longrightarrow C_j$ be the projection to the first coordinate. Then $pr_1(Interior(Im(\phi))) = C_j$.*

This Lemma implies that for each $c_j \in C_j$, there are open sets $U_{c_j}, V_{c_j} \in C_j$ such that $c_j \in U_{c_j}$ and $U_{c_j} \times V_{c_j} \subset Im(\phi)$. Since $\phi_1^* f = \phi_2^* f$, a.e., it follows that f is constant on U_{c_j}, a.e.. Since $pr_1(Interior(Im(\phi))) = C_j$ which is connected, f is independent of c_j.

Proof. Since the c_l's are all fixed for $l \neq j$, k is fixed. Since

$$R_{1,1} : K^2 \longrightarrow K',$$

$$R_{2,p} : K^{2(p-1)} \longrightarrow K'$$

are onto, one may choose regular values s, s' respectively of these two maps $R_{1,1}$ and $R_{2,p}$ such that $ss'S_{0,n}(c) = I$. Let $(g, h, c_j) \in M$ such that $R_{1,1}(g, h) = s$. Then (g, h, c_j) is a smooth point (The representation corresponding to (g, h, c_j) is irreducible) and the map ϕ_1 is regular at (g, h, c_j). It will then be sufficient to fix c_j and show that the map

$$\phi_2 : N(c_j) \longrightarrow C_j :$$

$$\phi_2(g, h) = (skc_jk^{-1})c_j(skc_jk^{-1})^{-1}$$

is regular at (g, h). Again, since both s, s' are regular values for the maps

$$R_{1,1} : K^2 \longrightarrow K',$$

$$R_{2,p} : K^{2(p-1)} \longrightarrow K',$$

s is a regular value for the map

$$R_{1,1} : N(c_j) \longrightarrow K'.$$

This implies that the differential $d\phi_2|_{(g,h)}$ is surjective. $\square$

Lemma 3.2 shows that f is independent of the variable c_j or the Γ_j-action is ergodic on M. If $F \in L^2(\mathrm{Hom}_C(\Sigma, K))$ is Γ-invariant, then again by the main result in [3], F is a characteristic function on C. By Lemma 3.2 and induction, F is a characteristic function on $\prod_{j=0}^{k} C_j$ for all $n \geq k \geq 0$. In particular, it is a characteristic function of C_0 (when $k = 0$). Theorem 1.1 follows.

References

[1] Atiyah, M., Bott, R., The Yang-Mills Equations Over Riemann Surfaces, *Philos. Trans. Roy. Soc. London*, Ser. A 308 (1982), 523-615.

[2] Goldman, W., The symplectic nature of fundamental groups of surfaces, *Adv. Math.* 54 (1984), 200-225.

[3] Pickrell, D., Xia, E., Ergodicity of Mapping Class Group Actions on Representation Varieties, I. Closed Surfaces, *Preprint.*

Department of Mathematics, University of Arizona, Tucson, AZ 85721 (*Pickrell*)

National Center for Theoretical Sciences, Third General Building, National Tsing Hua University, No. 101, Sec 2, Kuang Fu Road,, Hsinchu, Taiwan 30043, Taiwan R.O.C. (*Xia*)

E-mail address: `pickrell@math.arizona.edu` *(Pickrell)*, `xia@math.umass.edu` *(Xia)*